

TABLE 25. NUMBER OF DAYS FROM LAST SPRAY TO HARVEST
AND RESTRICTED ENTRY INTERVALS (hours) (cont.)**

Common name	Trade name	Time limit in days for use before harvest						REI(h)
		Apple	Pear	Cherry	Nectarine	Peach	Plum	
methomyl	Lannate	14	-	-	1 (J)	4	-	72-96 (L)
methoxyfenozide	Intrepid	14	14	-	7	7	7	4
metiram	Polyram	(L)	-	-	-	-	-	24
myclobutanil	Rally, Nova	14	-	0	0	0	0	24
naphthalene acetic acid	NAA	(F)	-	-	-	-	-	-
naphthylacetamide	Amid-Thin	(F)	-	-	-	-	-	-
napropamide	Devrinol	35	35	35	35	35	35	12
norflurazon	Solicam	60	60	60	60	60	60	24
novaluron	Rimon	14	-	-	-	-	-	12
OFM sprayable	CheckMate OFM-F	0	0	0	0	0	0	0
pheromone								
oryzalin	Surflan	0	0	0	0	0	0	12
oxamyl	Vydate	14	14	-	-	-	-	48
oxyfluorfen	Goal	(I)	(I)	(I)	(I)	(I)	(I)	24
paraquat	various Gramoxone	0	0	28	28	14	28	12
permethrin	Ambush, Perm-UP, Pounce	(B)	(C)	3	14	14	-	12
pendimethalin	Prowl H2O	60	60	60	60	60	60	24
phosmet	Imidan	7	7	7 (D)	14	14	7	72
prohexadione-calcium	Apogee	45	-	-	-	-	-	12
pronamide	Kerb	0	0	0	0	0	0	24
propargite	Omite	(P)	(P)	-	-	-	-	168
propiconazole	Orbit, Tilt, PropiMax	-	-	-	0 (R)	0 (R)	0 (R)	24
pyraclostrobin	Cabrio	-	-	0	-	-	-	12
pyraclostrobin	Pristine	0	0	0	0	0	0	12
+boscalid								
pyridaben	Nexter	25	7	300	7	7	7	12
pyrimethanil	Scala	72 (W)	72 (W)	- (W)	2 (W)	2 (W)	2 (W)	12
pyriproxyfen	Esteem	45	45	14	14	14	14	12
rimosulfuron	Matrix	14	14	14	14	14	14	4
sethoxydim	Poast	14	14	25	25	25	(P)	12
simazine	Princep	0	0	0	-	0	0	12
spinetoram	Delegate	7	7	7	1	14	7	4
spinosad	SpinTor, Entrust	7	-	7	(I)	14	7	4
spirodiclofen	Envidor	7	7	7	7	7	7	12
spirotetramat	Movento	7	7	7	7	7	7	24
stylet oil	JMS Stylet Oil	0	0	(L)	(L)	(L)	(L)	4
streptomycin	Agri-mycin 17	50	30	-	-	-	-	12
sulfur		0	0	0	0	0	0	24
tebuconazole	Elite	-	-	0	0	0	-	12
terbacil	Sinbar	60	-	-	-	60	-	12
thiacloprid	Calypso	30	30	-	-	-	-	12
thiamethoxam	Actara	(S)	(S)	14	14	14	14	12
trifloxystrobin	Flint, Gem	14	14	1	1	1	1	12
tebuconazole + trifloxystrobin	Adament	-	-	1	1	1	-	24
thiophanate-methyl	Topsin M	0	1	1	1	1	1	12
thiram		0	-	-	-	7	1	24
2,4-D	various (E)	14	14	40	40	40	40	48
triadimefon	Triadimefon	45 (N)	45 (N)	-	-	-	-	12
triflumizole	Procure	14	14	-	-	-	-	12
zeta-cypermethrin	Mustang Max	14	14	14	14	14	14	12
Ziram		14	14	14	14	14	-	48

(B) Apply no later than petal fall.

(C) Prebloom application.

(D) Do not use as a foliar spray on sweet cherries.

(E) Check herbicide label to determine which species can be treated with a specific formulation of 2,4 - D.

(F) Apply no later than 2 1/2 weeks after bloom for thinning. As a stop-drop spray, do not apply NAA within 2 days of harvest.

(I) Registered for dormant or delayed-dormant foliar application.

(J) SLN 24(c) label for Virginia and West Virginia.

(L) See label.

(N) Maximum 24 oz per acre per season.

(O) Flint for pome fruits, Gem for stone fruits.

(P) Non-bearing trees only.

(Q) Apply before growth starts in spring and in the fall after harvest.

(R) Apply a maximum of 2 sprays during the period beginning 3 weeks before harvest through the day of harvest (0 day PHI). Do not apply to prunes.

(S) 35 days if over 2.75 oz/A; 14 days for rates up to 2.75 oz/A.

(T) Postbloom handgun application to lower 4 ft of trunk for borer control

(U) See label. Re-entry interval may be more restrictive than days-to-harvest limitations.

(V) 21 days if over 2 lbs/A; 30 days for pick-your-own.

(W) Do not make more than 2 applications of a Group 9 fungicide (Scala or Vanguard) within 30 days of harvest.

** Days to harvest can vary according to the formulation and percentage of active ingredient used in some cases. Be sure to read the label carefully before applying any pesticide. Use this chart only as a guide.