

Denitrification Management

Zachary M. Easton, Assistant Professor and Extension Specialist, Biological Systems Engineering, Virginia Tech
Emily Lassiter, Graduate Research Assistant, Biological Systems Engineering, Virginia Tech

What Is Denitrification?

The only permanent removal of nitrogen (N) within both soils and aquatic environments is microbial *denitrification*^{*}, which reduces reactive nitrogen in the form of nitrate (NO₃⁻) to inert dinitrogen gas (N₂), which comprises approximately 78 percent of our atmosphere, or 4,000 trillion tons. Denitrification currently returns about half of excess (i.e., above plant requirements and potentially mobile) agricultural nitrogen sources to the atmosphere and could perhaps do more if effectively utilized within pollution-reduction strategies.

Soil microorganisms need oxygen to survive, and when the soil becomes saturated, oxygen is depleted, often referred to as *anaerobic*. Some *heterotrophic microbes* can get the oxygen needed to survive from nitrite (NO₂⁻) and NO₃⁻. When microbes utilize this oxygen, dinitrogen gas is created in a process called denitrification. Under certain conditions partial denitrification can occur, which results in the production of nitrous oxide (N₂O) or nitric oxide (NO), both powerful *greenhouse gases*. These gases are then lost to the atmosphere. In order to more fully understand what denitrification is and how it occurs, a brief explanation in the context of the nitrogen cycle is warranted.

The Nitrogen Cycle

Nitrogen is a naturally occurring element that is required by both plants and animals for growth and reproduction. As shown in figure 1, the nitrogen cycle can quickly become complex, with many nitrogen species interacting and constant transformations occurring under various conditions. The most abundant species of nitrogen — free, unreactive nitrogen in the atmo-

sphere, dinitrogen gas — is not in a form that is readily usable by plants or animals and does not combine readily with other elements because it is very stable. Thus, a substantial input of energy is required to break the N₂ bonds and create bioavailable nitrogen via nitrogen fixation. The nitrogen cycle consists of six well-understood steps and one that has been newly discovered that are discussed in greater detail below.

1. Nitrogen Fixation

Three processes are responsible for nitrogen fixation, which is the conversion of N₂ into a form that can be used by plants and animals (i.e., reactive or bioavailable nitrogen).

- **Atmospheric fixation** by lightning.
- **Biological fixation** by certain microbes (*Rhizobia* and *Bradyrhizobia* bacteria in root nodules), alone or in a symbiotic relationship with some plants such as legumes (e.g., soybeans, beans, peas). All biological nitrogen fixation requires the enzyme *nitrogenase*.
- **Industrial fixation** via the Haber-Bosch process used to produce fertilizer products such as urea that require a large amount of energy, primarily from non-renewable fossil fuels.

2. Immobilization

Abiotic reactions that sequester mineral nitrogen (e.g., ammonium fixation (*sorption*) by clay minerals) and *biotic* uptake and conversion of nitrogen to proteins and cell tissue by plants and microbes. Organisms mainly use nitrogen in its mineral forms (inorganic) NO₃⁻ and

* Terms defined in the glossary at the end of this publication are italicized the first time they appear in the text.

ammonium (NH_4^+), but it may also include uptake of soluble organic nitrogen. Biotic immobilization is also referred to as accumulation.

3. Mineralization

Microbial decomposition of organic nitrogen to mineral forms (reverse of immobilization). Ammonification is a subset of mineralization that results in the formation of NH_4^+ .

4. Ammonia Volatilization

Production of ammonia gas (NH_3) from NH_4^+ (as a dissolved ion) in soils with high concentrations of NH_4^+ and a pH greater than 7.5.

5. Nitrification

Stepwise microbial oxidation of NH_4 to NO_2^- to NO_3^- by the autotrophic bacteria *Nitrosomonas* and *Nitrobacter* respectively. Nitrifiers under anaerobic conditions can also produce the greenhouse gases NO and N_2O .

6. Denitrification

Microbially controlled reduction of nitrate to dinitrogen gas. In this reaction, organic carbon serves as an energy source, and nitrogen oxyanions (NO_3^- , NO_2^-) or oxides (NO, N_2O) serve as the terminal electron acceptor in the absence of free oxygen:



During each step organic carbon is utilized and CO_2 and H_2O are produced. Denitrification is the link completing the nitrogen cycle, converting reactive nitrogen (N_i) back to dinitrogen.

7. Anammox

Anaerobic oxidation of ammonium utilizing carbon dioxide (bicarbonate in aqueous environment) as an electron donor and nitrite as an electron acceptor. Anammox converts 90 percent of the reactant nitrogen to N_2 while converting 10 percent to NO_3^- . This process was discovered in a denitrifying wastewater treatment reactor in 1994. However, it is unclear how prevalent this process is in the natural environment.

In summary, the main forms of nitrogen present in the environment are nitrogen gas, organic N, ammonium/ammonia, and nitrate.




Figure 1. The nitrogen cycle, showing the forms of nitrogen and the mechanisms by which the various nitrogen species are transformed.

Environmental Impacts

While low N levels often limit biological production both in the terrestrial and marine environments, excess reactive nitrogen underlies a suite of environmental and health problems. Environmental degradation resulting from elevated nitrogen levels includes *eutrophication* of receiving water bodies, *global acidification*, stratospheric ozone depletion, and tropospheric accumulation of ozone and aerosols, which can lead to respiratory illness, cardiac disease, and cancer in humans.

Nitrogen can contaminate potable water. High concentrations of nitrate are toxic to infants — resulting in *methemoglobinemia* (blue baby syndrome) — as well as to livestock, while nitrite reacting with secondary amines forms carcinogenic nitrosamines.

Disruption of the global nitrogen cycle also has a multitude of potential impacts on the other major chemical cycles of carbon, phosphorus, and sulfur, which have become a topic of increasing study, particularly with respect to global climate change.

Thus, mitigating the effects of anthropogenic nitrogen inputs is vital to protection of our health and natural resources while continuing to produce food and energy for a growing population. Agricultural productivity is dependent on inputs of manufactured nitrogen fertilizer while energy production from fossil fuels also releases nitrogen that would not naturally be made available. Promoting and managing denitrification is paramount because the human population is sustained by anthropogenic increase in reactive nitrogen.




Figure 2. Various pathways that nitrogen can take in the environment (Rivett et al. 2008; reproduced with permission).

Figure 2 shows the various pathways that nitrogen can take in the environment. The fate of nitrogen depends on many conditions, including soil texture, organic matter content, soil moisture status, soil oxygen levels, soil pH, and temperature. Runoff and *leaching* are two, often large, pathways that transport nitrogen throughout the environment. Nitrogen runoff not only removes valuable plant nitrogen, but can also contribute to eutrophication of water bodies. Excessive nitrogen leaching also removes valuable plant nitrogen, can contaminate groundwater reservoirs and wells, and is very difficult to remove.

Incomplete denitrification or the production of harmful greenhouse gases N₂O and NO can degrade air quality. Nitrogen can be taken up by plants and microorganisms, stored in the soil (as organic nitrogen), or immobilized by sorption to soil.

Denitrification Management

Denitrification management (e.g., promoting conversion of excess NO₃⁻ to N₂) is thought to have some of the highest potential in agro-ecosystems. Globally, 75 percent of anthropogenic N_r is utilized in agro-ecosystems, of which approximately half is lost to the atmo-

sphere (NH₃, NO, N₂O, N₂) or to water (NO₃⁻); complete denitrification to N₂ accounts for only a minor portion.

Reduction of excess reactive nitrogen from a watershed scale to global scale must rely on multiple strategies employed in concert. There are four major management categories:

1. Minimize NO and N₂O emissions from fossil fuel combustion.
2. Improve nitrogen uptake efficiency in crops.
3. Increase nutrient management of livestock operations.
4. Expand access to sewage treatment globally.

Yet even these load reductions will not reduce reactive nitrogen inputs sufficiently to restore the balance in the global nitrogen cycle. The quantity of excess N_r as well as its persistence in the environment determines its impact because as nitrogen is transported through the environment and undergoes transformation, a single molecule may have multiple negative impacts, such as eutrophication (NO₃⁻), methemoglobinemia (NO₂), or as a greenhouse gas (N₂O, NO). This concept, known as the “nitrogen cascade,” emphasizes the importance of balancing anthropogenic nitrogen inputs with denitrification. Denitrification management — promoting reactive nitrogen removal throughout the cascade — is increasingly recognized as an essential component of managing the global nitrogen cycle, especially with respect to diffuse pollution.

Restoration and construction of wetlands and *riparian* buffers for nitrogen removal are examples of denitrification management applicable to a wide range of landscapes. However, many areas of our landscape are natural denitrification hot spots.

Denitrification occurs primarily under anaerobic conditions by heterotrophic microbes and, assuming sufficient NO₃⁻, denitrification rates are high in wet, carbon-rich soils.




Figure 3. Data from denitrification field study. The left figure indicates areas of the landscape likely to be saturated (i.e., anaerobic), with black dots representing sampling locations. The plot on the right is of denitrification rates from those sampling locations. Notice that the denitrification rates are substantially higher in the wet areas.

Figure 3 shows results from a field study of denitrification potential across a landscape. Notice in this figure that the highest denitrification rates are found in areas of the landscape that are most likely to be saturated (i.e., wet, often anaerobic or oxygen deficient). These denitrification hot spots also tend to be riparian areas with minimal disturbance, where soil-organic carbon can accumulate (remember, carbon is also a critical component in because it serves as the energy source to drive denitrification).

Limitations and Unknowns

Partial Denitrification

Under certain circumstances, harmful greenhouse gases such as NO and N₂O can be produced, and NO₂⁻ reacts to form a carcinogen. It is unclear exactly when this occurs.

Significant greenhouse gas production has been observed in other nitrogen removal systems utilizing denitrification, including constructed wetlands, which have been shown to act as both sources and sinks of nitrogen gases, occasionally producing high emissions of N₂O.

Increased temperature and certain carbon types — such as those with higher *labile* carbon content — can increase N₂O production during denitrification.

What You Can Do

- Protect areas of the landscape that act as natural denitrification hotspots. These would include frequently saturated areas, riparian areas, and wetlands.
- Plant residue is an important source of carbon and nitrogen for nitrification and denitrification. Increasing soil carbon content has the potential to provide in-field denitrification under certain conditions.
- Increase water residence time and contact with biological surfaces in ditch networks. Allowing some vegetation to grow, which slows water velocity and provides carbon, can do this.

- Fertilizer type can influence the magnitude of nitrification and denitrification and, as a result, nitrogen fluxes. For example, injected anhydrous ammonia commonly has higher NO production than other fertilizers.
- Recent work suggests that the use of controlled drainage in fields with subsurface tile drains can both prevent offsite transport of excess nitrogen and provide increased denitrification during periods of prolonged saturation.

Acknowledgements

We would like to thank David Sample, assistant professor and Virginia Cooperative Extension specialist, Virginia Tech; Joshua Faulkner, assistant professor and West Virginia University Extension specialist; and David Moore, Virginia Cooperative Extension agent and unit coordinator, Middlesex County, for their helpful and insightful reviews.

Additional Resources

Denitrification Research Coordination Network – www.denitrification.org

Food and Agriculture Organization of the United Nations (FAO), Natural Resources Management and

Environment Department, and the International Fertilizer Industry Association. 2001. "Factors Regulating Nitrous Oxide and Nitric Oxide Emission." Chapter 2 in *Global Estimates of Gaseous Emissions of NH₃, NO and N₂) From Agricultural Land*. Rome: FAO. www.fao.org/docrep/004/y2780e/y2780e02.htm

National Academy of Engineering Grand Challenges for Engineering – www.engineeringchallenges.org/cms/8996/9132.aspx

References

Rivett, M. O., S. R. Buss, P. Morgan, J. W. N. Smith, and C. D. Bement. 2008. "Nitrate Attenuation in Groundwater: A Review of Biogeochemical Controlling Processes." *Water Research*, 42(16): 4215-32.

Glossary of Terms

Abiotic – Nonliving factors that affect living organisms.

Anaerobic – Chemical reactions that proceed without the presence of oxygen.


Biotic – Consisting of living organisms.

Bradyrhizobia – See Rhizobia.

Denitrification – The transformation of nitrates to nitrites, nitric oxide, nitrous oxide, and ultimately, dinitrogen gas, which is lost to the atmosphere (i.e., the nitrogen cycle). Denitrification is fostered by anaerobic (see Anaerobic), carbon-rich conditions.

Eutrophication – Water body response to excess nutrients like nitrogen and phosphorus, generally resulting in excessive plant growth that often results in hypoxia — the depletion of oxygen in the water.

Global acidification – Also referred to as ocean acidification, it is the process of the current and ongoing decrease in the pH of the ocean. While increasing atmospheric carbon dioxide (CO₂) levels are the primary cause of ocean acidification, nitrogen oxides (N₂O, NO) also contribute substantially. The pH is reduced when these oxides are dissolved in seawater, which causes an increase in the hydrogen ion (H⁺) in solution (note that H⁺ is acidic and lowers the pH):



Greenhouse gases – A compound in the earth's atmosphere that blocks or traps heat that would normally radiate out of the atmosphere. The primary greenhouse gases are carbon dioxide, methane, nitrous oxide, and ozone.

Heterotrophic microbes – Microbes that obtain energy, carbon, and reducing equivalents for reactions from organic compounds.

Labile – The ability of a substance change easily. Labile carbon would mean carbon that is easily digested or utilized by microbes.

Leaching – The movement of water or dissolved constituents, usually downward; also referred to as percolation, out of the root zone.

Methemoglobinemia – Acute poisoning, primarily of infants, caused by nitrate ingested at high concentrations. Nitrate is converted to nitrite in an infant's digestive system, and the nitrite reacts with hemoglobin (that carries oxygen to all parts of the body) to form methemoglobin, which does not carry oxygen. The level of oxygen being carried throughout the body decreases as hemoglobin is converted to methemoglobin. As the oxygen level decreases, the baby is suffocated.

Microbial decomposition – The breakdown of compounds or organic matter into smaller compounds by microorganisms.

Nitrobacter – A bacteria that converts nitrite into nitrate.

Nitrogenase – Enzymes used to fix atmospheric nitrogen gas (N₂) to convert to ammonia (NH₃).

Nitrosomonas – A bacteria that converts ammonia into nitrite.

Rhizobia – Soil microbes that fix nitrogen once established in legumes. The microbes maintain a symbiotic (i.e., both microbe and plant benefit) relationship with the legume. The microbes supply ammonia or amino acids to the plant and, in return, receive organic acids as a carbon and energy source.

Riparian – Describes the near stream area, often forested, and subject to periodic saturation.

Sorption – The physical bonding of one substance to another.